



## Effect of CO<sub>2</sub> elevation on Shisham growth at nursery stage

Nirmal<sup>1\*</sup> • Meena Bakshi<sup>2</sup> • Hukum Singh<sup>2</sup>

<sup>1</sup> Indian Institute of Farming System Research, Modipuram

<sup>2</sup> Forest Research Institute, Dehradun

### ARTICLE INFO

### ABSTRACT

#### Article history:

Received: 17 December, 2021

Revision: 24 December, 2021

Accepted: 30 December, 2021

Key words: Shisham, CO<sub>2</sub> elevation, growth, nursery stage.

The effects of elevated carbon dioxide concentration (CO<sub>2</sub>) were evaluated on Shisham at nursery stage for height, diameter, number of branches and number of leaves by raising them for a year in open top chamber (OTC) with six levels of CO concentrations viz. Control, 400ppm, 600ppm, 800ppm, 1000ppm and 1200ppm. Morphological studies associated with it would be beneficial in understanding the overall mechanism underlying growth and development of Shisham in response to elevated CO<sub>2</sub>. Stem height was observed to be maximum at 1000 ppm CO<sub>2</sub> Concentration (85.84±4.67) whereas diameter (10.21±0.22), number of leaves (56.82±4.74) and number of branches (6.53±0.50) were recorded maximum in control.

### 1. Introduction

Shisham (*Dalbergia sissoo* Roxb.) is an excellent timber species worldwide occurring naturally in many Asian countries like India, Afganistan, Pakistan, Bangladesh, Bhutan, Burma and Nepal. It is widely adapted to varied edaphic and climatic conditions, so it is one of the most preferred species for afforestation and reforestation programmes. Farmers of North India also plant this species in agroforestry systems for profitable economic returns and is well accepted for social forestry programmes.

Research has shown that plants respond positively at elevated CO<sub>2</sub> (Amthor 1995) which is the raw material of photosynthesis and has great influence on plant physiology, growth, structure and function of plant species. The current atmospheric CO<sub>2</sub> concentration is about 380  $\mu\text{mol mol}^{-1}$ , which is far below the optimum concentration of plant photosynthesis. Generally, plants grown at elevated CO<sub>2</sub> relative to those grown at ambient CO<sub>2</sub> often exhibit increased growth, and photosynthesis with improved water use efficiency. The effects of elevated CO<sub>2</sub> are manifested by changes in photosynthesis. Studies of both deciduous and evergreen plants have shown that elevated CO<sub>2</sub> leads to increased photosynthesis (Ells worth *et al.* 2004) and decreased water conductance (Calfapietra *et al.* 2005). Although much is known regarding the effects of elevated CO<sub>2</sub> on agricultural crops but forest species including

*Dalbergia sissoo* Roxb. an important industrial and agroforestry tree species, has received much less attention. Studies on effect of CO<sub>2</sub> elevation would be useful to understand the response of this economically important agroforestry tree species in purview of changing climate.

### 2. Materials and Methods

The present study was conducted in the (OTC) Open Top Chamber facility at the campus of Forest Research Institute, Dehradun, India which is situated at Latitude 30°20'10.31" N, Longitude 77°59'55.32" E and altitude of 650 amsl. The study was conducted during the period March 2018 to February 2019. Clones of Shisham (Clone 14, 232 and 86) were raised *via* cuttings in OTCs. To maintain the homogeneity of the experiment the cuttings were raised in equal size pots. 18 plants per CO<sub>2</sub> concentration were taken in five chambers each.

Commercial grade 100% CO<sub>2</sub> gas was supplied to the chambers through CO<sub>2</sub> gas cylinder and maintained at set levels (400 ppm, 600 ppm, 800 ppm, 1000 ppm and 1200 ppm) using gas regulators, pressure pipelines, solenoid valves, rotameters, sampler, pump, CO<sub>2</sub> analyzer, PC linked Program Logic Control (PLC) and Supervisory Control and Data Acquisition (SCADA). The uniformity of CO<sub>2</sub> gas was maintained by diluting CO<sub>2</sub> gas with air by a 120 liter capacity air compressor.

\*Corresponding author: [nirmal@icar.gov.in](mailto:nirmal@icar.gov.in)

### Growth attributes/ Biometric traits

Each plant's height (cm), collar diameter (mm), number of leaves, number of branches were recorded seasonally viz. spring, summer, rainy, and winter at interval of four months throughout the period of the entire experiment.

#### Height (in cm)

The height of emerged plants from root collar to tip of the shoot were recorded seasonally at interval of four months with the help of meter scale in centimeter (cm) seasonally.

#### Collar diameter (in mm)

The collar diameter (diameter at root collar) was recorded using digital caliper in millimeter seasonally at interval of four months.

#### Number of leaves and branches

Leaf number and branches were counted seasonally at interval of four months. Being pinnately compound for counting each set of leaflets emerging from the branch was taken as a single leaf.



### 3. Results and Discussion


Primary results indicated that altered CO<sub>2</sub> concentration had significant effects on growth parameters of Shisham at nursery stage. Maximum height was observed in Shisham growing at 1000 ppm of CO<sub>2</sub> (85.84±4.67cm) followed by Control (70.66±3.44 cm) and 800 ppm (68.69±3.32 cm) respectively. Among the different levels of CO<sub>2</sub> concentration maximum collar diameter was observed in Control (10.21±0.22 mm) followed by 1000 ppm (9.64±0.24 mm) and 800 ppm (9.36±0.25 mm). Maximum number of branches was seen in control (6.53±0.50) followed by 1000 ppm (4.58±0.45) and 800 ppm (4.18±0.40) respectively. Maximum number of leaves were recorded in Control (56.82±4.74) followed by 1000 ppm (43.15±4.28) and 1200 ppm (37.61±3.74) of CO<sub>2</sub> concentration respectively (Table 1).

CO<sub>2</sub> enrichment has shown varying effect on agricultural and horticultural crops. Usui *et al.* (2014) found that heat tolerant cultivars of rice showed higher grain percentage along with maintaining grain quality in Free Air CO<sub>2</sub> Enrichment (FACE) experiment studies.

**Table 1.** Effect of CO<sub>2</sub> elevation on growth parameters of *Dalbergia sissoo* at nursery stage

CO <sub>2</sub> Concentration	Height (cm)	Diameter (mm)	No. of Branches	No. of Leaves
Control	70.66 <sup>ab</sup> ±3.44	10.21 <sup>b</sup> ±0.22	6.53 <sup>b</sup> ±0.50	56.82 <sup>b</sup> ±4.74
400	68.05 <sup>ab</sup> ±3.43	8.22 <sup>a</sup> ±0.25	2.81 <sup>a</sup> ±0.25	28.76 <sup>a</sup> ±2.09
600	63.91 <sup>a</sup> ±2.87	9.21 <sup>ab</sup> ±0.25	3.43 <sup>a</sup> ±0.36	27.71 <sup>a</sup> ±2.45
800	68.69 <sup>ab</sup> ±3.32	9.36 <sup>ab</sup> ±0.25	4.18 <sup>a</sup> ±0.40	31.90 <sup>a</sup> ±2.24
1000	85.84 <sup>b</sup> ±4.67	9.64 <sup>ab</sup> ±0.24	4.58 <sup>ab</sup> ±0.45	43.15 <sup>ab</sup> ±4.28
1200	63.85 <sup>a</sup> ±3.34	8.10 <sup>a</sup> ±0.23	3.85 <sup>a</sup> ±0.45	37.61 <sup>a</sup> ±3.74
F value	3.314	4.373	6.278	7.749
p value	0.01	0.001	0	0
S/NS	S	S	S	S





**Fig 1.** Effect of CO<sub>2</sub> elevation on growth parameters of *Dalbergia sissoo* at nursery stage



**Fig 2.** *Dalbergia sissoo* plants in Open top chambers (OTC) at different CO<sub>2</sub> level concentrations; a. Control, b. 400ppm, c. 600ppm, d. 800ppm, e. 1000ppm and f. 1200ppm.

Kimball *et al.* (2007) showed that enrichment at +300 ppm CO<sub>2</sub> above ambient (ambient was about 350 ppm in 2007) resulted in an overall growth (fruit + wood) stimulation of 70% ( $\pm$  12%) after 17 years in sour orange (*Citrus aurantium* L.) trees. Madan *et al.* (2012) concluded that elevated CO<sub>2</sub> increase yield in rice varieties but high temperature reduces grain quality. Chakraborty *et al.* (2015) showed that CO<sub>2</sub> enrichment resulted in significant increase in growth, leaf area and dry matter production in *Brassica* cultivars. Tomato grown under enhanced CO<sub>2</sub> concentrations were found to flower earlier and produce 30% more marketable fruit than ambient air by Hickleton and Jolliffe (1978).

Increased CO<sub>2</sub> levels have also been observed to stimulate growth in forest species. In an assessment, modern oaks growing at an average CO<sub>2</sub> of 330 ppm, growth sensitivity to temperature was found about twice than that of paleo oaks growing at an average CO<sub>2</sub> of 230 ppm by Voelker *et al.* (2017). In their studies Purohit and Habibi (2010) found that among certain tree species; *Acharus sapota*, *Wrightia tomentosa*, *Feronia limonia*, *Terminalia bellerica* and *Celastrus paniculatus* cultural growth, survival percentage, overall growth parameters showed significant enhancement over the control under elevated CO<sub>2</sub>. Large synergistic gains from higher CO<sub>2</sub> and nutrients was detected with nutrients added in forest experiments on maturing pine by Oren *et al.* (2001). Smith *et al.* (2013) found that the mean effect of CO<sub>2</sub> enrichment on aboveground woody biomass was +29, +22 and +16% for *A. glutinosa*, *F. sylvatica* and *B. pendula*, respectively in monoculture. Dawes *et al.* (2011) depicted larger growth in *Larix* growing under elevated CO<sub>2</sub> but not in *Pinus*. Density of algae (*Trebouxia* sp.) was found significantly higher after 380 days exposure to the CO<sub>2</sub>-enriched environment by Ismail *et al.* (2017). In another important observation Mohan *et al.* (2007) noticed that shade tolerant species showed increment whereas shade intolerant species didn't survive better with CO<sub>2</sub> enrichment.

#### 4. Conclusion

Previous research has shown a positive response of plants to elevated CO<sub>2</sub> (Amthor 1995) in our primary results it has been clearly depicted that CO<sub>2</sub> elevation had significant increase in height of Shisham, collar diameter, number of branches and number of leaves when compared to ambient (Fig. 1). Shisham was found to outperform in terms of growth at ambient levels (400ppm) of CO<sub>2</sub> in controlled environment, optimum growth was seen at 1000ppm which declined steeply thereafter at 1200ppm which might be the tolerance threshold for the species.

#### 5. References

- Amthor JS (1995). Terrestrial higher-plant response to increasing atmospheric [CO<sub>2</sub>] in relation to the global carbon cycle. *Glob Change Biol.* 1(4): 243-274
- Calfapietra C, Bernacchi CJ, Centritto M, Sharkey TD (2005). Photosynthetic responses to increased CO<sub>2</sub> and air pollutants. *Terrestrial Photosynthesis in a Changing Environment: A Molecular, Physiological and Ecological Approach*, ed. Flexas J, Loreto F, Medrano H, Cambridge University Press.
- Chakraborty K, Uprety DC, Bhaduri D, (2015). Growth, Physiology and Biochemical Responses of Two Different Brassica Species to Elevated CO<sub>2</sub>. *Proc. Natl. Acad. Sci., India, Sect. B Biol. Sci.* doi 10.1007/s40011-015-0615-9
- Dawes MA, Hattenschwiler S, Bebi P, Hagedorn F, Handa IT, Korner C, Rixen C (2011). Species-specific tree growth responses to 9 years of CO<sub>2</sub> enrichment at the alpine treeline. *Journal of Ecology*, 99: 383–394
- Ellsworth DS, Reich PB, Naumburg ES, Koch GW, Kubiske ME, Smith SD (2004). Photosynthesis, carboxylation and leaf nitrogen responses of 16 species to elevated CO<sub>2</sub> across four free-air CO<sub>2</sub> enrichment experiments in forest, grassland and desert. *Global Change Biol.* 10:2121–2138.
- Hickleton PR, and Jolliffe PA, (1978). Effects of greenhouse CO<sub>2</sub>, enrichment on the yield and photosynthetic physiology of tomato plants. *Can. J. Plant. Sci.* 58: 801-817.
- Ismail A, Marzuki SD, Yusof NBM, Buyong F, Said MNM, Sigh HR and Zulkifli ARZ (2017). Epiphytic Terrestrial Algae (*Trebouxia* sp.) as a Biomarker Using the Free-Air-Carbon Dioxide-Enrichment (FACE) System. *Biology* 2017, 6-19; doi: 10.3390/biology6010019
- Kimball BA, Idso SB, Johnson S, Rillig MC (2007). Seventeen years of carbon dioxide enrichment of sour orange trees: final results. *Global Change Biology* 13: 2171-2183.
- Madan P, Jagadish SVK, Craufurd PQ, Fitzgerald M, Lafarge T, Wheeler TR (2012). Effect of elevated CO<sub>2</sub> and high temperature on seed-set and grain quality of rice. *Journal of Experimental Botany*, 63(10): pp. 3843–3852
- Mohan JE, Clark JS, Schlesinger WH (2007). Long-Term CO<sub>2</sub> Enrichment of a Forest Ecosystem: Implications for Forest Regeneration and Succession. *Ecological Applications*, 17(4), pp. 1198-1212

- Oren R, Ellsworth DS, Johnsen KH, Phillips N, Ewers BE, Maier C, Schafer KVR, McCarthy H, Hendrey G, McNulty SG, Katul GG (2001). Soil fertility limits carbon sequestration by forest ecosystems in a CO<sub>2</sub>-enriched atmosphere. *Nature* Vol. 411: 469-472
- Purohit SD, Habibi N (2010). Effect of CO<sub>2</sub> Enrichment on In Vitro Growth, Hardening and Acclimatization during Micropropagation of Some Tree Species of Aravallis in Rajasthan Proc. IVth IS on Acclim. and Establ. of Micropropagated Plants Ed.: J. Prakash Acta Hort. 865, ISHS 2010
- Smith AR, Lukac M, Hood R, Healey JR, Douglas L, Godbold DL (2013). Elevated CO<sub>2</sub> enrichment induces a differential biomass response in a mixed species temperate forest plantation. *New Phytologist* 198: 156–168. doi: 10.1111/nph.12136
- Usui Y, Sakai H, Tokida T, Nakamura H, Nakagawa H, Toshihiro (2014). Hasegawa Heat-tolerant rice cultivars retain grain appearance quality under free-air CO<sub>2</sub> enrichment *Rice*: 7:6. doi: 10.1186/s12284-014-0006-5
- Voelker SL, Stambaugh MC, Brooks JR, Meinzer FC, Lachenbruch B, Guyette RP (2017). Evidence that higher [CO<sub>2</sub>] increases tree growth sensitivity to temperature: a comparison of modern and paleo oaks. *Oecologia. Global Change Biology*. doi: 10/1007/s00442-17-3831-6.